

The phase diagram of the anisotropic Spin-1 Heisenberg Chain

A. L. Malvezzi and F. C. Alcaraz
Departamento de Física
Universidade Federal de São Carlos
13565-905, São Carlos, SP, Brasil

KEY WORDS: Haldane gap, exact diagonalization,
conformal field theories, density matrix renormalization group

In 1983, Haldane conjectured that the isotropic antiferromagnetic spin-S Heisenberg chain has a gap or not depending if the spin is an integer or half-integer[1]. This conjecture is today well established [2, 3]. In the case where an exchange z-anisotropy is present the situation is less clear and the location of the point where the massive Haldane phase appears is still controversial even for the spin 1 case. The Hamiltonian in this last case is the spin-1 XXZ chain

$$\mathcal{H} = \sum_{j=1}^L \left(S_j^x S_{j+1}^x + S_j^y S_{j+1}^y + \lambda S_j^z S_{j+1}^z \right) \quad (1)$$

where S_j^x , S_j^y and S_j^z are the spin 1 operators.

From finite-size studies (for lattices sizes up to $L = 14$) and by exploring the results of conformal invariance[4] Alcaraz and Moreo[5] conjectured that for $\lambda < \lambda_c = 0$ the Hamiltonian (1) is in a massless X-Y phase with critical fluctuations governed by a conformal field theory with central charge $c = 1$. Moreover in the critical phase ($-1 \leq \lambda \leq 0$) the dimensions of operators $x_{n,m}^{(b)}$, $x_n^{(s)}$ governing the correlation functions in the bulk and in the surface are given by[5]

$$x_{n,m}^{(b)} = n^2 x_p + \frac{m^2}{4x_p}, \quad x_n^{(s)} = 2x_p n^2 \quad (2)$$

where

$$x_p = \frac{\pi - \cos^{-1}(\lambda)}{4\pi}, \quad n, m = 0, \pm 1, \pm 2, \dots \quad (3)$$

The Haldane phase is expected to appear when the operator $x_{0,1}^{(b)} = \frac{1}{4x_p} = \frac{\pi}{\pi - \cos^{-1}(\lambda)}$ becomes relevant, which according to eqs. (2) and (3) occurs at $\lambda = \lambda_c = 0$. This fact explains[5] some exact degeneracies appearing in the spectra of (1) with free ends and is corroborated by the results of ref. [6].

More recently Yajima and Takahashi[7] by calculating the spectra of (1) for lattice sizes up to $L = 16$ concluded that the Haldane phase should starts at $\lambda = \lambda_c = 0.068 \pm 0.003$. Since this result destroy the conjecture (2) we decide to make a more precise calculation of λ_c by using the density matrix renormalization group (DMRG) introduced by White[8], which enable us to make spectral calculations in much larger lattice sizes. The direct way to estimate λ_c would be by a direct mass gap evaluation of (1) in a periodic chain. However our results shows that for $\lambda \approx 0$ the finite-size gaps are very small (the transition at λ_c should be of Kosterlitz-Thouless type) and it is

very difficult to decide if the point $\lambda = 0.065$ (the worse estimative against the conjecture (2)) is in a massive phase or not. Another way to estimate λ_c is by calculating the exponents η_x , governing the correlation $\langle S_i^x S_{i+r}^x(r) \rangle \sim r^{-\eta_x}$. The critical phase disappears when η_x reaches the value[7, 9] $2x_{1,0}^{(b)} = 1/4$ or the surface exponent[5] $x_1^{(s)}$ reaches the value $1/4$.

Since the DMRG is much more precise for spectral calculations in open chains we calculate $x_1^{(s)}$ in order to estimate λ_c . This is done by using the finite-size relations coming from the conformal invariance of the infinite system[4]. The exponent $x_{(1)}^s$ is obtained by extrapolating ($L \rightarrow \infty$) the sequence

$$x_1^{(s)}(L) = \frac{E_1^1(\lambda, L) - E_1^0(\lambda, L)}{E_2^0(\lambda, L) - E_1^0(\lambda, L)}. \quad (4)$$

where $E_j^n(\lambda, L)$ is the j^{th} energy in the sector where $\sum S_i^z = n$ of the Hamiltonian (1) with free ends.

In table 1 we show for the special cases $\lambda = 0$ and $\lambda = 0.065$ the estimated values for $x_1^{(s)}(L)$ for lattices up to $L = 48$. In our DMRG calculations we keep $k = 50$ and 80 states in the truncated Hilbert space[8]. A good estimator for the errors is $1 - P_k$, where P_k is the trace of the truncated density matrix. The values in table 1 were obtained by taking $P_k \rightarrow 1$.

At $\lambda = \lambda_c$, the operator $x_{0,1}^{(b)}$, which corrects the finite size scaling[5], should be marginal and logarithmic corrections in the sequence (4) are expected. In fig. 1 we show the data of table 1, at $\lambda = 0$, together with a mean square fit in the form

$$x_1^{(s)}(L) = \frac{1}{4} + a (\ln L)^b. \quad (5)$$

showing a good numerical fit. A similar polynomial fit, which would be the case if $\lambda_c > 0$, although possible shows larger deviations. On the other hand the data of table 1 at $\lambda = 0.065$ are not well fitted by (5). As we can see from table 1 the data at $\lambda = 0.065$ clear indicate a limiting value $x_1^{(s)}(\infty)$ bigger than $1/4$. In order to see this more clearly we calculated at $\lambda = 0.065$ this estimator for $L = 60$ sites and obtain $x_1^{(s)}(60) = 0.25023$, which indicates that $\lambda = 0.065$ is already a point in the Haldane phase.

In conclusion, our results indicate that, in agreement with the conjecture (2), the Haldane phase appears for $\lambda > \lambda_c = 0$, and the logarithmic corrections appearing at this point explains the numerical controversial in earlier numerical calculations.

Acknowledgments

This work was supported in part by CNPq and Fundação de Amparo à Pesquisa do Estado de São Paulo - FAPESP, SP-BRASIL.

References

- [1] F. D. M. Haldane: Bull. Am. Phys. Soc. **27** (1983) 464; Phys. Rev. Lett. **50** (1983) 1153.
- [2] M. P. Nightingale and H. W. Blöte: Phys. Rev. B **33** (1986) 659; M. Takahashi: Phys. Rev. B **38** (1988) 5188, Phys. Rev. Lett. **62** (1989) 2313; K. Nomura: Phys. Rev. B **40** (1989) 2421; T. Sakai and M. Takahashi: Phys. Rev. B **42** (1990) 1090.
- [3] R. Botet and R. Jullien: Phys. Rev. B **27** (1983) 613; R. Botet R. Julien and M. Kolb: Phys. Rev. B **28** (1983) 3914; S. R. White and D. A. Huse: Phys. Rev. B **48** (1993) 3844.
- [4] J. L. Cardy: Nucl. Phys. B **270** [FS16] (1986) 186; J. L. Cardy: *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic, New York, 1986), Vol. 11.
- [5] F. C. Alcaraz and A. Moreo: Phys. Rev. B **46** (1992) 2896.
- [6] F. C. Alcaraz and Y. Hatsugai: Phys. Rev. B **46** (1992) 13914.
- [7] M. Yajima and M. Takahashi: J. Phys. Soc. Jpn. **63** (1994) 3634.
- [8] S. R. White: Phys. Rev. Lett. **69** (1992) 2863; Phys. Rev. B **48** (1993) 10345.
- [9] H. J. Schultz: Phys. Rev. B **34** (1986) 6372.

Figure Caption

Fig. 1 - The data of table 1 together with the mean square fit (continuum curve).

Table Caption

Table 1 - Finite size estimatives of x_1^s , obtained from eqs. (4) for $\lambda = 0$ and $\lambda = 0.065$.

Table 1

L	$\lambda = 0.0$	$\lambda = 0.065$
8	0.236462774	.246639580
16	0.237501420	.247661670
24	0.237964476	.248398065
32	0.238271248	.248954979
40	0.238500913	.249401545
48	0.238685182	.249774106